

Characteristics of hydrothermal convection in inclined layers: implications for hydrothermal activity at slow-spreading axis

F. Fontaine, M. Cannat, J. Escartin, D. Dusanur

Institut de Physique du Globe- Géosciences Marines, Boite 89, 4 place Jussieu., Paris, 75252 France

The thermal structure of segments along (slow-spreading) mid-ocean ridges is likely to be a key parameter controlling the distribution, dynamics and geometry of hydrothermal systems. It is usually considered that the depth of penetration of hydrothermal fluids at the ridge axis is a function of the depth to the brittle-ductile transition. At slow-spreading axis, it is likely that this depth varies both along- and across-axis, with a deepening of several kilometers from the segment center towards its ends [e.g., Hooft et al., 2000]. This geometry is a consequence of focused melt supply to the segment center, resulting in the episodic and localized injection of magma bodies in the crust, as observed at the Lucky Strike segment of the Mid-Atlantic ridge [Singh et al., 2005]. In order to study the effect of such slopes of the basal temperature on the dynamics of slow-spreading axis hydrothermal systems, we ran a series of two-dimensional numerical models of hydrothermal convection. As a first approximation and following previous studies [e.g., Rabinowicz et al., 1999], we assume that these systems can be represented as rectangular and inclined permeable layers. The models are single-phase and incorporate realistic fluid properties and permeabilities. We have explored the cases of slopes ranging from 0 to 15°, aspect ratios from 1 to 16, and permeabilities up to 10^{-14} m². The basal slope controls the number of convective cells. As the slope increases, the ratio of the size of the downflow and upflow areas increases. Above a critical slope the circulation is uni-cellular and composed of a broad recharge zone and a focused discharge zone, and encompassing the whole length of the segment. We will present the implication of our models for the distribution of vent sites along slow-spreading ridge segments. The segment-scale circulation and focused outflow obtained could also explain the elevated heat flux at some of the main sites found along slow-spreading ridges like Rainbow and TAG [German and Lin, 2004].