

A positive magnetic anomaly at Rainbow hydrothermal site in ultramafic environment

Jérôme DYMENT¹, Kensaku TAMAKI², H  l  ne HOREN³, Yves FOUQUET⁴, Kaori NAKASE⁵, Michiko YAMAMOTO⁵, Morgane RAVILLY⁶, Mitsuko KITAZAWA^{1,7}

1: CNRS UMR 7097 Laboratoire de G  osciences Marines, Institut de Physique du Globe de Paris, 4 place Jussieu 75005 Paris, France, 2: University of Tokyo, Department of Geosystem Engineering, University of Tokyo, Yayoi 1-1-1, Bunkyo-ku, Tokyo 113-0032, Japan, 3: CNRS UMR 8538 Laboratoire de G  ologie, Ecole Normale Sup  rieure, 24 rue Lhomond, 75005 Paris, France, 4: Laboratoire de G  ochimie et M  talog  nie, DRO/GM, IFREMER Centre de Brest, BP 70, 29280 Plouzan  , France, 5: Formerly at Ocean Research Institute, University of Tokyo, Minamidai 1-15-1, Nakano-ku, Tokyo, Japan, 6: CNRS UMR 6538 Domaines oc  aniques, Intitut Universitaire Europ  en de la Mer, Universit   de Bretagne Occidentale, 1 place N. Copernic, 29280 Plouzan  , France, 7: Ocean Hemisphere Research Center, Earthquake Research Institute, University of Tokyo, Yayoi 1-1-1, Bunkyo-ku, Tokyo 113-0032, Japan

Most of the hydrothermal vent systems on basaltic substratum present a negative magnetic anomaly, i.e. a deficit of magnetization, related to either thermal demagnetization associated to the high temperature at the vents, or to alteration of the basalt titanomagnetite to less magnetized titanomaghemite, or both. Our investigations shows that hydrothermal vent systems on ultramafic substratum may be associated to strong positive anomalies instead, which origin is currently investigated.

In May 2001, hydrothermal site Rainbow (MAR 36  14'N) was explored through a variety of experiments using IFREMER deep-sea ROV Victor, including one long dive fully devoted to a Deep Sea Three Components Magnetometer (DSTCM) survey. A total of 22 lines, amounting for more than 16 km, were collected, among which 19 lines, 700-800 km long and 60 m apart, covered quite densely the whole site area. The magnetic data were acquired about 1 to 10 m above seafloor, providing an ultra-high resolution never achieved so far on a hydrothermal system in ultramafic environment.

Both the magnetic effects from the ROV and the main geomagnetic field were evaluated and removed. The absolute magnetization of the seafloor was estimated in the following way: (1) synthetic magnetic anomalies were calculated along the ROV tracks, assuming a local 2D topography (as given by the immersion and altitude of the ROV) and a unit (1A/m) magnetization of the seafloor, (2) observed and synthetic anomalies were compared on sliding windows, and if the coherence between both signals is high, an estimate of the magnetization is given by the ratio between observed and modeled anomalies. Surprisingly, site Rainbow exhibits a very strong magnetization (higher

than 28 A/m). Although this observation can be explained either by the formation of magnetite during the serpentinisation process or by formation of highly magnetic minerals within the sulfurs in relation to the peculiar settings of the site, preliminary rock magnetic analysis of representative samples suggests that the major magnetic bearer is magnetite located within sulfurs impregnating the altered ultramafic rocks.

A cruise proposal has been submitted to further document the magnetic anomaly (by carrying out a denser magnetic survey of the site) and the magnetic bearer (by collecting samples for further rock magnetic experiment). The proposal has been ranked high and may be scheduled in 2007.